

Floquet topological semimetal with nodal helix:

“The Stark effect in topological semimetals”

Kun Woo Kim

Institute for Basic Science

Quantum transport:

1. Transport of disordered boundary modes.
2. TI nanowire's magneto-conductance.
3. Metal-to-Insulator transition via disorder.

Floquet quantum matter:

1. Photocurrent in surface Dirac fermion.
2. Weyl semimetal under strong E-field.
3. Floquet Nodal helix semimetal.

Disordered Floquet quantum matter:

1. Anomalous Floquet Anderson insulator.
2. Quantum kicked rotor in 4D.
3. Quantum walk with chiral symmetry in $!D$.


Collaborators:

Gil Refael (Caltech) group.
Junho Seo (KRISS) group.
Hee Chul Park (IBS) group.


Naoto Nagaosa (RIKEN) group.
Kwon Park (KIAS) group.

Alexander Altland (Cologne) group.




Floquet topological insulators



Oka and Aoki, PRB 79 081406 (2009)



Lindner, Refael, Galitski, Nature Physics, 7, 490 (2011)




Runder et al. PRX 3, 031005 (2013)

Outline of this talk

1. The Stark effect and Bloch oscillation in solids: theory and experiments
2. Wannier-Stark ladders in topological semimetals:
 - **Weyl semimetal + E-field:** [KWK](#), WR Lee, YB Kim, K Park, Nat. comm. **7**, 13489
 - **Nodal helix semimetal + E-field:** [KWK](#), HW Kwon, K Park, arXiv:1808.04079

The Stark effect



Consider a hydrogen atom. Due to the rotational symmetry of Coulomb interaction with proton, electronic energy levels are degenerated.

The application of electric field breaks the inversion symmetry and energy level splitting increases with the field strength. (discovered in 1913, Nobel prize in 1919).

Along with the Zeeman effect, the experimental observation of the Stark effect provided the confirmation of perturbation theory in quantum mechanics.

Wannier-Stark ladder from the Schrodinger equation



In 1960 G.H. Wannier predicted the Stark effect for a Bloch electron in a constant electric field.

$$\left[-\frac{1}{2m} \frac{\partial^2}{\partial x^2} + V(x) + eEx \right] \psi(x) = \epsilon \psi(x)$$

If “ ϵ ” is an eigenvalue of the Schrodinger equation with eigenfunction $\psi(x)$, there is a series of ladder like eigenvalues “ $\epsilon + maeE$ ” with eigenfunction $\psi(x - ma)$:

$$\left[-\frac{1}{2m} \frac{\partial^2}{\partial x^2} + V(x) + eE(x - ma) \right] \psi(x - ma) = \epsilon \psi(x - ma)$$

The Stark effect of electrons in lattice: Wannier-Stark ladder

Interestingly, Wannier's solution from the abelian approximation naturally contains the Zak phase:


$$\epsilon = \left(m + \frac{1}{2\pi} \gamma^{\text{Zak}} \right) aeE + \frac{a}{2\pi} \int_0^{2\pi/a} \epsilon(k) dk$$

Integer m , lattice constant a ,
a dispersion relation $\epsilon(k)$.



$$\gamma^{\text{Zak}} = \int_0^{2\pi/a} \left\langle u_k \left| i \frac{\partial}{\partial k} u_k \right. \right\rangle dk$$

Considering $i \frac{\partial}{\partial k} = \hat{x}$ as a position operator, the Zak phase is a measure of Bloch electron's position within unit cell, or a polarization. It is then naturally translated to a potential energy under an external electric field.

➔ Shifting of intra-cell position



The Stark effect : perspective in momentum space



Under an electric field, a free electron will be accelerated indefinitely. $\hbar \frac{dk}{dt} = eE$


On the other hand, electrons under a periodic potential will oscillate in momentum and real space following a dispersion relation, $\epsilon(k) \rightarrow \epsilon(k - eEt/\hbar)$.

To travel across the Brillouin zone n-times: $T = \frac{2\pi n}{a} \frac{\hbar}{eE}$.

A Bloch electron shouldn't be scattered during the travel: $T < \tau_s \quad \rightarrow \quad n \frac{\hbar}{\tau_s} < eaE$

Experimental limitation: a lattice unit cell of 5\AA , E-field 10 (kV/cm), $v_F = 10^5$ (m/s) yields the length of Bloch oscillation: ~ 1 (μm), which is one order larger than a typical mean free path 100 (nm).

Wannier-Stark ladder in experiments: semiconductor superlattice




A semiconductor superlattice structure is used to simulate artificial semiconductor with much larger lattice spacing ($\sim 100\text{\AA}$). With electric field strength ($\sim 10\text{kV/cm}$), the Wannier-Stark ladder was observed.

Wannier-Stark ladder in experiments: semiconductor superlattice

Wannier-Stark ladder in experiments: semiconductor superlattice

A semiconductor superlattice structure is used to simulate artificial semiconductor with much larger lattice spacing ($\sim 100\text{\AA}$). With electric field strength ($\sim 10\text{kV/cm}$), the Wannier-Stark ladder was observed.


For a given E-field strength, the number of ladder indicates the localization length of the Bloch oscillation. The stronger the E-field, the more localized is the Bloch electron.



A semiconductor superlattice structure is used to simulate artificial semiconductor with much larger lattice spacing ($\sim 100\text{\AA}$). With electric field strength ($\sim 10\text{kV/cm}$), the Wannier-Stark ladder was observed.

For a given E-field strength, the number of ladder indicates the localization length of the Bloch oscillation. The stronger the E-field, the more localized is the Bloch electron.

Wannier-Stark ladder in experiments: cold atoms in optical lattice



Standing wave potential with gravity to realize the Stark Hamiltonian:

$$V(x) = \cos(kx) + mgx.$$

Two counter propagating waves

$$\begin{aligned} V(x, t) &= \cos(w_1 t + k_1 x) + \cos(w_2 t - k_2 x), \\ &= 2 \cos\left(\bar{w}t + \frac{1}{2}\Delta kx\right) \cos\left(\frac{1}{2}\Delta w t + \bar{k}x\right), \\ &\sim \cos\left(\bar{k}\left[x - \frac{1}{2}at^2\right]\right). \end{aligned}$$

The linearly increasing phase difference, $\Delta w = w_2 - w_1 = at\bar{k}$, is simulating the gravity and it thus provides an external “electric” field.


Outline of this talk

1. The Stark effect and Bloch oscillation in solids: theory and experiments
2. Wannier-Stark ladders in topological semimetals:
 - **Weyl semimetal + E-field:** [KWK](#), WR Lee, YB Kim, K Park, Nat. comm. 7, 13489
 - **Nodal helix semimetal + E-field:** [KWK](#), HW Kwon, K Park, arXiv:1808.04079


Our working regime: $\frac{h}{\tau_s} < eaE < \text{Bandwidth}$

Weyl semimetal and the Zak phase

Weyl node is located in 3-dimensional space. We can think of the system as a collection of 1-dimensional wires along z-axis in momentum space, $\vec{k}_\perp = (k_x, k_y)$. For each wire, we can compute the Zak phase $\gamma^{Zak}(\vec{k}_\perp)$.



$$\gamma^{Zak} = \int_0^{2\pi/a} \langle u_{\vec{k}} | i \frac{\partial}{\partial k_z} u_{\vec{k}} \rangle dk_z$$





Wannier-Stark ladder in Weyl semimetal

Momentum resolved energy spectrum is:

$$\epsilon(\vec{k}_\perp) = \left(m + \frac{1}{2\pi} \gamma^{\text{Zak}}(\vec{k}_\perp) \right) aeE + \frac{a}{2\pi} \int_0^{2\pi/a} \epsilon(k_z, \vec{k}_\perp) dk_z \quad \text{with integer } m.$$

The zero energy cut of spectrum will contain a open line segment from the Zak phase winding.




Outline of this talk


1. The Stark effect and Bloch oscillation in solids: theory and experiments
2. Wannier-Stark ladders in topological semimetals:
 - **Weyl semimetal + E-field:** [KWK](#), WR Lee, YB Kim, K Park, Nat. comm. 7, 13489
 - **Nodal helix semimetal + E-field:** [KWK](#), HW Kwon, K Park, arXiv:1808.04079

Our working regime: $\frac{h}{\tau_s} < eaE < \text{Bandwidth}$

Nodal-line semimetal and nodal helix in time domain



Nodal-line



Nodal-helix

Nodal-line semimetal is characterized by a Zak phase of a loop enclosing a nodal line.

$$\gamma^{\text{Zak}} = \oint \langle u_{\vec{k}} | i \frac{\partial}{\partial \vec{k}} u_{\vec{k}} \rangle d\vec{k} = \pi$$

Consider a gapless 2D Dirac fermion under circularly polarized light. The trace of Dirac point will make a helix in momentum-time domain.

$$H(\vec{k}, t) = [k_x - A \cos(\Omega t)] \sigma_x + [k_y - A \sin(\Omega t)] \sigma_y$$

From a Floquet Hamiltonian to a Stark Hamiltonian

Technical side

Start with a continuum Dirac Hamiltonian: $H(\vec{k}, t) = [k_x - A\cos(\Omega t)] \sigma_x + [k_y - A\sin(\Omega t)] \sigma_y$

Because $H(t) = H(t+T)$, eigenvectors assume the following form from the Bloch theorem:

$$\psi(t) = e^{-i\epsilon t} \phi(t) \quad \text{with} \quad \phi(t) = \phi(t + T),$$

where a time-periodic function can be Fourier decomposed: $\phi(t) = \sum_n e^{-in\Omega t} \tilde{\phi}_n$.

Writing the Schrodinger equation in the Floquet basis, $\tilde{\phi}_n$:


$$i \frac{\partial}{\partial t} \psi(t) = H(\vec{k}, t) \psi(t) \quad \longrightarrow \quad \sum_m [\tilde{H}_{nm} - n\Omega \delta_{nm}] \tilde{\phi}_m = \epsilon \tilde{\phi}_n$$

Where $\tilde{H}_{nm} = \frac{1}{T} \int_0^T H(t) e^{-i(n-m)\Omega t} dt$. The Floquet operator can be thought of as a hopping between Floquet site n and m with a static electric field with strength Ω .

We ended up having 3-dimensional system with E-field from 2-dimensional time-periodic Hamiltonian.

Nodal helix in graphene

With two gapless Dirac fermion in graphene, two nodal helices are generated.




In the high driving frequency limit, this is well known Floquet quantum spin Hall system by Kitagawa et al. We work this problem in the low frequency limit to preserve the 3-dimensionality, and a strong intensity of light regime to induce a sizable ring of helix in momentum space.

Nodal helix in graphene: Zak phase and LDOS at zero energy


A reminder of Wannier-Stark ladder:

$$\epsilon(\vec{k}_\perp) = \left(m + \frac{1}{2\pi} \gamma^{\text{Zak}}(\vec{k}_\perp) \right) aeE + \frac{a}{2\pi} \int_0^{2\pi/a} \epsilon(k_\parallel, \vec{k}_\perp) dk_\parallel$$

- a. Zak phase $\gamma^{\text{Zak}}(\vec{k}_\perp)$
- b. Momentum resolved LDOS at $E=0$.
- c. LDOS($E=0$) from the abelian approximation.
- d. Magnified view of **b**.




Nodal helix in graphene: dispersion relation



Energy dispersion relation from the abelian approximation shows the Zak phase π shift across the ring marked by yellow shade. By the inter-band tunneling, the discontinuity in spectrum is smoothly connected with energy gap opening.

Here the color code indicates the fraction of eigenfunction belonging to the (conduction/valence) solution from the abelian approximation.


Nodal helix in graphene: dispersion relation



The energy gap is exponentially suppressed with the reduction of driving frequency Ω . In this sense, we argue that the gapless nature is approximately preserved in the Wannier-Stark ladder with an external ‘electric’ field. Thus, the observation of nodal helix semimetal carrying Zak phase shift π will be possible.

$$\mathcal{E}_{gap} \cong \frac{\Omega A}{\sqrt{2\pi k(A+k)}} e^{-\frac{2(A-k)^4}{k(A+k)\Omega^2}}$$

Experiment: irradiated 2D Dirac surface state in 3D topological insulator



Parameters in this experiment:


Driving frequency = 0.2 (eV)

E-field strength = 3.3×10^5 (V/cm)

Size of ring $k_0 = 0.0165$ ($1/\text{\AA}$)

We want to point out that this experiment can be interpreted as a series of Wannier-Stark ladder with the Zak phase π change reflected in the spectrum.

A single 2D gapless Dirac fermion: intermediate driving frequency regime



Even with moderate regime of driving frequency, we can see the Wannier-Stark ladder carries a significant amount of conduction (red)/valence (blue) band characters from the abelian approximation.

1st column: Wannier-Stark ladder from the Abelian approximation.

2nd column: Wannier-Stark ladder without approximation.

3rd and 4th column: realistic situation. We limit the number of accessible Floquet bands.

Summary and Outlook

1. The Bloch oscillation and the Stark effect in lattice are reviewed. The Zak phase is picked up by the Bloch oscillation, and it appears in spectral property.
2. Weyl semimetal shows the winding of the Zak phase around projected Weyl nodes. Thus, the Bulk spectrum shows a open Fermi arc.
3. A time periodic 2-dimensional system can be mapped into a 3-dimensional system under a static external electric field. The circular motion of the Dirac node realizes a helix nodal semimetal.
4. The time evolution of Bloch wave-packet in WSL semimetal through Landau-Zener tunneling is something to look at.

Thank you for your attention.

Technical slide

A few line of algebra can solve the problem:
$$\left[-\frac{1}{2m} \frac{\partial^2}{\partial x^2} + V(x) + eEx \right] \psi(x) = \epsilon \psi(x)$$

Rewrite the equation in the basis of Bloch wave function $\psi(x) = \sum_{nk} B_n(k) \psi_{nk}(x)$:

$$\left[-\frac{1}{2m} \frac{\partial^2}{\partial x^2} + V(x) + eEx \right] \sum_{nk} B_n(k) \psi_{nk}(x) = \epsilon \sum_{nk} B_n(k) \psi_{nk}(x),$$

Then, take an inner product by an Bloch wave function from the left:



$$\left[\epsilon_n(k) + eEi \frac{\partial}{\partial k} + \sum_m eE \left\langle u_{mk} \left| i \frac{\partial}{\partial k} u_{nk} \right. \right\rangle \right] B_n(k) = \epsilon B_n(k)$$

By taking one band only, We can obtain:
$$B_n(k) = \exp \left(-\frac{i}{eE} \int_0^k \left[\epsilon - \epsilon_n(k') - eE \left\langle u_{nk'} \left| i \frac{\partial}{\partial k'} u_{nk'} \right. \right\rangle \right] dk' \right)$$


From the periodic condition, energy quantization:
$$\epsilon = maeE + \frac{a}{2\pi} \int_0^{2\pi/a} \left[\epsilon_n(k') + eE \left\langle u_{nk'} \left| i \frac{\partial}{\partial k'} u_{nk'} \right. \right\rangle \right] dk'$$

Weyl semimetal in experiments

1. Topological insulator multilayer proposal (Burkov and Balents, PRL **107**, 127205)





(a) $m=0$



(b) $m \neq 0$

(Δ_S : coupling within TI. Δ_D : coupling between TI's. m : Zeeman coupling on TI surface.)

2. Cold atoms in optical lattice (Gross and Bloch, Science **357**, 995)



$|\uparrow\rangle$ and $|\downarrow\rangle$

ill-defined boundary and a limited transport measurement.

Wannier-Stark ladder to characterize Weyl semimetallic phase.

From the Zak phase to the first Chern number

The Zak phase $\gamma^{\text{Zak}}(\vec{k}_{\perp})$ changes by 2π around projected Weyl node in momentum space $\vec{k}_{\perp} = (k_x, k_y)$.

This is because, the change of the Zak phase is the first Chern number:

$$\begin{aligned} \oint \left[\frac{\partial}{\partial \vec{k}_{\perp}} \gamma^{\text{Zak}}(\vec{k}_{\perp}) \right] d\vec{k}_{\perp} &= \oint \frac{\partial}{\partial \vec{k}_{\perp}} \left[\int_0^{2\pi/a} \langle u_{\vec{k}} \left| i \frac{\partial}{\partial k_z} u_{\vec{k}} \right\rangle dk_z \right] d\vec{k}_{\perp} \\ &= \oint \int_0^{2\pi/a} \left[\left\langle \frac{\partial}{\partial \vec{k}_{\perp}} u_{\vec{k}} \left| i \frac{\partial}{\partial k_z} u_{\vec{k}} \right\rangle - \left\langle \frac{\partial}{\partial k_z} u_{\vec{k}} \left| i \frac{\partial}{\partial \vec{k}_{\perp}} u_{\vec{k}} \right\rangle \right] dk_z d\vec{k}_{\perp} \\ &= \text{Winding number from a torus to a sphere.} \end{aligned}$$

A Weyl semimetal can be viewed as a collection of 2D Chern insulators in momentum space. And the Fermi arc surface state is a collection of chiral edge states.